



This article is available to subscribers. [Subscribe now](#). Already have an account? [Sign in](#)

ORIGINAL ARTICLE

FREE PREVIEW

ARCHIVE

# Measles Outbreak in a Fully Immunized Secondary-School Population

Tracy L. Gustafson, M.D., Alan W. Lievens, M.D., Philip A. Brunell, M.D., Ronald G. Moellenberg, B.S., Christopher M.G. Buttery, M.D., and Lynne M. Sebulster, Ph.D.[et al.](#)

March 26, 1987

N Engl J Med 1987; 316:771-774

DOI: 10.1056/NEJM198703263161303

## Abstract

An outbreak of measles occurred among adolescents in Corpus Christi, Texas, in the spring of 1985, even though vaccination requirements for school attendance had been thoroughly enforced. Serum samples from 1806 students at two secondary schools were obtained eight days after the onset of the first case. Only 4.1 percent of these students (74 of 1806) lacked detectable antibody to measles according to enzymelinked immunosorbent assay, and more than 99 percent had records of vaccination with live measles vaccine. Stratified analysis showed that the number of doses of vaccine received was the most important predictor of antibody response. Ninety-five percent confidence intervals of seronegative rates were 0 to 3.3 percent for students who had received two prior doses of vaccine, as compared with 3.6 to 6.8 percent for students who had received only a single dose. After the survey, none of the 1732 seropositive students contracted measles. Fourteen of 74 seronegative students, all of whom had been vaccinated, contracted measles. In addition, three seronegative students seroconverted without experiencing any symptoms.

We conclude that outbreaks of measles can occur in secondary schools, even when more than 99 percent of the students have been vaccinated and more than 95 percent are immune. (N Engl J Med 1987; 316:771-4.)

This article is available to subscribers.

SUBSCRIBE

OR

PURCHASE ARTICLE →

---

---

## Funding and Disclosures



We are indebted to Ray Falk, Melinda Gonzales, Gerry Heitkamp, and America Gonzales of the Corpus Christi School District; to Helen Reeves, Nancy Clary, and Carl Johnson of the health department; to Dr. Joe Oshman, Dr. Robert Bosquez, Dr. Patrick Brosnen, Mr. Wheeler Lipes, and all the other health professionals in both Corpus Christi and the Texas Department of Health for their help with the serologic survey; and to Judy Schwitzer for secretarial support.

---

## Author Affiliations



From the Division of Infectious Diseases Epidemiology and the Bureaus of Communicable Disease Service and Epidemiology, Texas Department of Health, Austin; the Department of Pediatrics, University of Texas Health Science Center, San Antonio; and the Corpus Christi–Nueces County Health Department, Corpus Christi, Tex. Address reprint requests to Dr. Gustafson at Infection Control and Prevention Analysts, 1122 N. Alma #234, Richardson, TX 75081.